

8. Verbitskiy E. V., Dinastiya E. M., Baranova A. A. *et al.* // Dyes and Pigments. 2018. Vol. 159. P. 35–44.
9. Verbitskiy E. V., Baranova A. A., Khokhlov K. O. *et al.* // Chemistry of Heterocyclic Compounds. 2018. Vol. 54. P. 604–611.
10. Kvashnin Y. A., Kazin N. A., Verbitskiy E. V. *et al.* // ARKIVOC. 2016. Vol. V. P. 279–300.
11. Verbitskiy E. V., Kvashnin Y. A., Baranova A. A. *et al.* // Russ. Chem. Bull. 2018. Vol. 67. P. 1078–1082.
12. Verbitskiy E. V., Kvashnin Y. A., Baranova A. A. *et al.* // Dyes and Pigments. 2019. Vol. 168. P. 248–256.

** This work was supported by the Russian Foundation for Basic Research (Research Project № 18-29-23045 МК).*

УДК 547.892.4

E. Y. Zelina, T. A. Nevolina, M. G. Uchuskin


*Perm State University,
614990, Russia, Perm, Bukireva St., 15,
zelina.e@psu.ru*

THE PAAL-KNORR REACTION IN THE SYNTHESIS OF 1,2-ANNULATED PYRROLES*

Keywords: Furan, Paal-Knorr synthesis, 1,4-Diketone, Domino reaction.

An original effective protocol for the synthesis of 1,2-annulated pyrroles has been developed. This new process features the simultaneous formation of both pyrrole and diazepine cores as a result of intramolecular Paal-Knorr reaction of 1,4-dicarbonyl compounds forming during an acid-catalyzed hydrolysis of the corresponding furans **1**. A wide variety of pyrrolo[1,2-*a*][1,4]diazepines **3** and pyrrolo[1,2-*d*][1,4]diazepines **4**, prospective pharmacological agents, was synthesized (*path a*) [1]. In addition to this, a simple and effective methods for the synthesis of substituted (het)arene-annulated pyrrolo[1,2-*d*][1,4]diazepines **5** and pyrrolo[1,2-*a*]quinoxalines **6**. The developed approach is based on the acid-catalyzed hydrolysis of substituted furans **2** with the formation of the corresponding 1,4-diketones followed by the key reductive cyclization with the formation of a broad range of nitrogen heterocycles **5**, **6** (*path b*) [2, 3].

The optimal reaction conditions were found, the scope and limitations of the developed method were defined. Cytotoxicity of synthesized 1,2-annulated pyrroles against cell lines HEK293T, VA13, MCF7, A549 and MCF10A was investigated. Research results will be presented in the report.



References

1. Zelina E. Y., Nevolina T. A., Sorotskaja L. N. et al. // J. Org. Chem. 2018. Vol. 83. P. 11747–11757.
2. Zelina E. Y., Nevolina T. A., Skvortsov D. A. et al. // J. Org. Chem. 2019. Vol. 84. P. 13707–13720.
3. Zelina E. Y., Nevolina T. A., Sorotskaja L. N. et al. // Tetrahedron Lett. 2020. Vol. 61. In the Press.

* This work was supported by Russian foundation for basic research (project № 19-43-590007 p_a).

УДК 547.786.1

Д. А. Аксенов, Н. А. Аксенов,
Е. В. Александрова, А. В. Аксенов

Северо-Кавказский федеральный университет,
355009, Россия, г. Ставрополь, ул. Пушкина, 1а,
Mr.twos@mail.ru

ONE-POT ПОЛУЧЕНИЕ 2-(3-ОКСОИНДОЛИН-2-ИЛ)-2-АРИЛАЦЕТОНИТРИЛОВ РЕАКЦИЕЙ ИНДОЛОВ С НИТРОАЛКЕНАМИ*

Ключевые слова: индолы, нитроалкены, фосфористая кислота, нитрилы.

Наша группа заинтересована в разработке каскадных гетероциклизаций нитросоединений с помощью кислот Бренстеда. Нами было показано, что нитроалкены могут быть использованы в качестве синтетических эквивалентов